WHAT ARE **OPIOIDS?**

Opioids are a class of drugs that reduce pain and promote feelings of pleasure and relaxation. Taking too many opioids slows your breathing which leads to lack of oxygen in your brain, coma, heart stopping, and death. This process can be fast, so time is critical to saving a life.

Common opioids include prescription opioids, heroin, and illicit fentanyl.

WHAT ARE THE SIGNS OF AN **OVERDOSE?**

- Blue lips/fingernails (ashen if darker skin)
- Breathing that is infrequent, uneven or has stopped
- Deep snoring or gurgling noises
- Unresponsive to pain stimulus
- Pinpoint pupils

WHAT IS NALOXONE?

- Naloxone (Narcan,[®] Evzio[®]) is a drug that temporarily reverses the dangerous effects of an opioid overdose
- Naloxone can be given every 2-3 minutes, but only works if opioids are in your body
- Works for reversing fentanyl overdoses
- A person cannot get high on it or become addicted to it

EFFECTIVE ONLY FOR 30 TO 120 MINUTES

resources

OVERDOSE ACTION NALOXONE TRAINING: overdoseaction.org For opioid overdose naloxone training

TREATMENT CENTERS

HOME OF NEW VISION homeofnewvision.org 734.975.1602 - Even Month Birthdays

DAWN FARMS
dawnfarm.org
734.669.8265 - Odd Month Birthdays

If you live in:

Livingston County, MI - 517.546.4126 Monroe County, MI - 734.243.7340 Lenawee County, MI - 517.263.8905

HARM REDUCTION AND SYRINGE ACCESS

UNIFIED miunified.org 734.961.1082

Funding was provided by Michigan Institute for Clinical & Health Research UL1TR000433, Michigan Department of Health and Human Services and Community Mental Health Partnership of Southeast Michigan

© 2017 The Regents of the University of Michigan

All rights reserved.

opioid safety & naloxone

TAKE A.C.T.I.O.N.
AND SAVE A LIFE
www.overdoseaction.org

Opioid Overdose <u>Deaths are Pre</u>ventable.


If you suspect someone is experiencing an opioid overdose, have your naloxone accessible and take immediate A-C-T-I-O-N

HOW TO PREVENT OVERDOSE AND REDUCE HARM

- Do not mix with other drugs especially alcohol,benzodiazepines, and anything that makes you sleepy
- Store your medication out of reach of children or pets
- Reduce substance amounts
- Be aware of the tolerance changes in your body
- Go slow if injecting drugs
- Alternate injection sites
- Train family/friends to call for help and know where naloxone is stored and how to use it
- Go slow, never use alone, and carry naloxone
- Dispose unused medications properly: michigan.gov/doyourpart

INSTRUCTIONS FOR

intramuscular naloxone



- 1. Pop the orange flip top from the naloxone vial
- 2. Insert 23-25G, 1-1.5 inch needle into the vial and draw 1 mL of naloxone into syringe
- 3. Inject needle straight into muscle (through clothes, if needed), then push in plunger all the way

REMEMBER TO TAKE A.C.T.I.O.N



1: AROUSE (3 "S")

Shout their name

Shake shoulders vigorously in attempts to arouse

Sternal rub



2: CHECK FOR SIGNS OF OVERDOSE

Slowed or no breathing Blue/gray lips or fingernails Deep snoring/ Gurgling noises Unresponsive to pain



3: TELEPHONE 911 Call 911



4: INTRAMUSCULAR/ INTRANASAL NALOXONE

Give naloxone



5: OXYGEN

Do rescue breaths: 2 breaths initially, then 1 breath every 5-6 second OR Start CPR if you know how OR follow dispatch instructions



6: NALOXONE AGAIN

Repeat naloxone in 2-3 minutes if the person does not resume breathing.
Recovery position (side lying) if you must leave the person unattended or the person vomits.

Stay with the person if safe to do so until help arrives.



narcan nasalspray

Narcan® does not require assembly.







* Do NOT push the plunger until the nozzle is in the nostril. NO TEST SPRAY is needed. If still not breathing in 2-3 minutes, use the 2nd spray.

WHAT NOT TO DO IN OVERDOSE

Do not DELAY calling 911

Do not put the person in a bath or in shower

Do not give the person anything to drink

Do not inject the person with anything

Do not pour water over the face or slap too hard

Do not leave the person till help arrives